

## BIROn - Birkbeck Institutional Research Online

Tavacioglu, E.E. and Azanon Gracia, Elena and Longo, Matthew R. (2019) Perceptual distortions of 3-D finger size. *Perception* 48 (8), pp. 668-684. ISSN 0301-0066.

Downloaded from: <https://eprints.bbk.ac.uk/id/eprint/27597/>

*Usage Guidelines:*

Please refer to usage guidelines at <https://eprints.bbk.ac.uk/policies.html>  
contact [lib-eprints@bbk.ac.uk](mailto:lib-eprints@bbk.ac.uk).

or alternatively

RUNNING HEAD: Perception of 3-D Finger

## Perceptual Distortions of 3-D Finger Size

Ebru Ecem Tavacioglu<sup>1,2,3</sup>, Elena Azañón<sup>1,4,5,6</sup>, and Matthew R. Longo<sup>1</sup>

<sup>1</sup>Department of Psychological Sciences, Birkbeck, University of London, UK

<sup>2</sup>Department of Psychology, Ludwig-Maximilians-Universität München, Munich,  
Germany

<sup>3</sup>Psychology Department, Istanbul Sehir University, Turkey

<sup>4</sup>Institute of Psychology, Otto von Guericke University Magdeburg, Germany

<sup>5</sup>Center for Behavioral Brain Sciences, Otto von Guericke University Magdeburg,  
Germany

<sup>6</sup>Department of Behavioral Neurology, Leibniz Institute for Neurobiology, Magdeburg,  
Germany

Address correspondence to:

Matthew Longo

Department of Psychological Sciences

Birkbeck, University of London

Malet Street, London WC1E 7HX

United Kingdom

[m.longo@bbk.ac.uk](mailto:m.longo@bbk.ac.uk)



## Abstract

Our body is a volumetric, 3-dimensional object in the world, and we experience it as such. Existing methods for measuring the perceptual body image, however, have been based on judgments of 1-dimensional length or 2-dimensional images. We developed a new approach to the 3-D perceptual body image of the fingers by asking people to judge whether each finger would fit through rings of varying diameter. This task requires participants to conceptualize their finger as a volumetric object entering the ring. In two experiments, we used an adaptive staircase procedure to estimate the perceived size of each finger. There were systematic distortions of perceived 3-D finger size, with the size of index finger and (to a lesser extent) the middle finger underestimated. These distortions were unaffected by changes in hand posture. Notably, the pattern of distortions is qualitatively different from that found in previous research investigating 1-D finger length, suggesting that 3-D judgments of the body may differ in fundamental ways from 1-D judgments of individual body dimensions.

Keywords: Body Image, Fingers, Posture, Body Size

Distortions of perceived body size and shape are a core aspect of eating disorders such as anorexia nervosa (Bruch, 1962). A large literature has used body size estimation tasks to compare eating disordered patients and controls (e.g., Ben-Tovim, Whitehead, & Crisp, 1979; Garner, Garfinkel, Stancer, & Moldofsky, 1976; Horne, Van Vactor, & Emerson, 1991; Slade & Russell, 1973), and meta-analyses of this literature have consistently found that patients overestimate body width compared to controls (Cash & Deagle, 1997, Mölbert et al., 2017; Sepúlveda, Botella, & León, 2002; Smeets, Smit, Panhuysen, & Ingleby, 1997). Such differences, however, do not imply that healthy individuals show veridical judgments, and indeed many studies have reported systematic misperception of body size and shape in non-clinical samples (e.g., D'Amour & Harris, 2017; Dolan, Birtchnell, & Lacey, 1987; Dolce, Thompson, Register, & Spana, 1987; Fuentes, Longo, & Haggard, 2013; Halmi, Goldberg, & Cunningham, 1977; Hundleby & Bourguoin, 1993; Linkenauger et al., 2015, 2017; Sadibolova, Ferrè, Linkenauger, & Longo, 2019; for reviews see Thompson, 1986; Longo, 2017a). Such results suggest that far from being a sure sign of disease, distorted representations of the body may be a ubiquitous part of ordinary cognitive life.

The clinical literature on body size estimation has focused largely on the body parts such as the torso, waist, and hips, given their obvious importance to body image concerns in the context of eating disorders. A more recent literature has focused on investigating perceptual representations of the hand, a body part of particular interest given its central role in skilled action (Jeannerod, 1997). A range of distortions of hand representation have been identified, including hand representations involved in position sense (e.g., Longo & Haggard, 2010; Longo, Long, & Haggard, 2012; Saulton, Dodds, Bulthöff, & de la Rosa, 2015; Coelho, Zaninelli, & Gonzalez, 2017), tactile distance perception (e.g., Green, 1982; Taylor-Clarke, Jacobsen, & Haggard, 2004; Longo &

Haggard, 2011), tactile localization (e.g., Trojan et al., 2006; Mancini, Longo, Iannetti, & Haggard, 2011), and judgments of the configuration of landmarks within the hand (e.g., Longo, 2015a; Margolis & Longo, 2015).

Of particular relevance to the present study, three recent studies have used a simple perceptual matching task to measure the perceived length of the fingers (Longo & Haggard, 2012a; Longo, Mattioni, & Ganea, 2015b; Tamè, Bumpus, Linkenauger, & Longo, 2017a). In this task, participants were asked to compare the perceived length of each finger (i.e., the distance from the knuckle to the outstretched fingertip) to the length of a line shown on a computer monitor. The perceived length of each finger was estimated using either a staircase procedure (Longo & Haggard, 2012a; Longo et al., 2015b) or the method of adjustment (Tamè et al., 2017a). In each case, there was substantial underestimation of finger length, which showed a clear gradient across the hand. Underestimation was smallest for the thumb and increased progressively across the hand towards the little finger. Intriguingly, these distortions are qualitatively similar to, but smaller in magnitude than, distortions of proprioceptive hand maps, where participants point to the location of the knuckles and finger tips of their occluded hand (Longo & Haggard, 2010), and differences between fingers in both tactile sensitivity (Duncan & Boynton, 2007; Manser-Smith, Tamè, & Longo, 2018; Sathian & Zangaladze, 1996; Vega-Bermudez & Johnson, 2001) and cortical magnification factors (Duncan & Boynton, 2007).

These studies have described distortions of the hand on a single dimension (e.g., length). However, our body is a 3-dimensional (3-D), volumetric object, and we experience it as such. The 3-D nature of the body, in general, has not been well captured by existing methods of measuring the perceptual body image. The present study addresses this gap and investigates the body image of the fingers in 3-D. One class of

methods measures 1-D length, such as the moving caliper method (Slade & Russell, 1973), the adjustable light beam method (Thompson & Spana, 1988), the body image detection device (Ruff & Barrios, 1986), and the line length task described in the previous paragraph. Another class of methods measures 2-D proportions (i.e., aspect ratio), such as the distorting picture method (Glucksman & Hirsch, 1969), the body-distorting mirror (Traub & Orbach, 1964), the video distortion method (Allebeck, Hallberg, & Espmark, 1976), and the template matching task (Gandevia & Phegan, 1999). Some other studies have attempted to address this issue by obtaining estimates of body depth (e.g., Button, Fransella, & Slade, 1977; Halmi et al., 1977; Casper et al., 1979; Dolce et al., 1987), showing body images in profile (e.g., Freeman, Thomas, Solyom, & Hunter, 1984; Lindholm & Wilson, 1988; Brodie, Slade, & Rose, 1989; Fernández, Probst, Meermann, & Vandereycken, 1994; Urdapilleta, Aspavlo, Masse, & Docteur, 2010), or obtaining 1-D measures of perceived limb circumference (Horne et al., 1991; Salbach et al., 2007; Schneider et al., 2009a, 2009b; Keizer, van Elburg, Helms, & Dijkerman, 2016; Mölbert et al., 2016). None of these approaches, however, involve judgments about the body as a 3-D object, but about different dimensions of the body, individually. It is unclear whether the perceived shape of a body part in 3-D is fundamentally different from what might be inferred from measuring judgments of each dimension in isolation. Thus, the nature of distortions in 3-D space for the body, and in particular for the hand, remains unclear.

In the present study, we use a new approach to investigate the body image of the fingers in 3-D, by asking participants to make judgements about whether their fingers would fit into circular rings of differing diameters. Rather than judging the size of each finger in any single dimension, this task requires that the participant conceptualize their finger as a volumetric object entering and moving through the ring. Because the thickest

part of the finger is not necessarily at the base of the finger, this task also requires that participants consider the movement of the entire finger through the ring, and not merely judge cross-sectional area of any specific finger location. We used an adaptive staircase procedure to estimate the perceived size of each finger.

## Experiment 1

### *Method*

#### *Participants*

Twenty individuals (9 women, mean age = 32.4 years, range: 21-48 years) participated. Participants were all right-handed as assessed by the Edinburgh Handedness Inventory (Oldfield, 1971; mean = 85.6, range = 50 – 100). Participants reported no known abnormalities of tactile perception and normal or corrected-to-normal visual acuity. Participants provided written informed consent and were paid for their participation. Procedures were approved by the Department of Psychological Sciences Research Ethics Committee at Birkbeck.

#### *Stimuli*

Stimuli were 33 metal ring gauges (Digiflex, London, UK), examples of which are shown in Figure 1. The metal band of each ring was 2.35 mm in thickness. The inner diameter of each ring ranged from 12.1 to 23.4 mm in approximately equal steps.





**Figure 1:** Examples of the ring gauges used as stimuli. On each trial, the participant was shown one of the rings and judged whether one of their fingers would be able to fit through the ring. The penny is shown for scale.

### *Procedure*

Participants were seated at a table facing the experimenter with their hands placed on their lap. They were asked not to look at their hands, which were covered by a black cloth. On each block, the participant was asked to make judgments about the size of one of the five fingers of their left hand. On each trial, the experimenter placed one of the ring gauges in front of the participant at approximately the height of the participant's eyes. The participant was given the follow description of the task: "In this task, we will ask you to imagine that one of the fingers of your left hand is going through a ring placed in front of you. On each block, I will let you know which finger you should judge. Your task is to judge whether the ring is big enough that your finger could pass through it without any friction. That is, it would be fine if your finger touched the sides of the ring, so long as it didn't put any pressure on your skin. If it seems like the ring is big enough that your finger would fit through it without friction, say "yes". If it seems like the ring is not big enough, say "no".

The ring was placed on a custom-made plasticine support allowing it to rest upright giving the participant a clear view of its entire circumference at a viewing distance of approximately 50 cm. Responses were made verbally and were unspeeded.

The experiment consisted of 5 blocks, one for each finger, in random order. Each block consisted of 36 trials, for a total of 180 trials. On each trial, the QUEST Bayesian adaptive staircase algorithm (Watson & Pelli, 1983) as implemented in the Psychtoolbox (Brainard, 1997) for MATLAB (Mathworks, Natick, MA), was used to select which ring gauge to present. QUEST analyses the participant's history of previous responses to find the stimulus (i.e., the diameter of ring gauge) which will be most informative for estimating the true threshold (i.e., the stimulus size for which participants would be equally likely to say "yes" and "no"). On each trial, the size of ring gauge to be displayed was determined by QUEST and displayed to the experimenter on a monitor. Each block consisted of two interleaved QUEST staircases, each consisting of 18 trials, one starting with a large stimulus (26 mm diameter) and the other starting with a small stimulus (18.5 mm diameter). At the start of the experiment each participant completed five practice trials using a single staircase, and starting with the 18-mm diameter ring. The participant was allowed to take a short break midway through and after each block, but was not allowed to look at their hands at any point.

At the end of the experiment, the actual size of each finger was measured using the same set of ring gauges by identifying the smallest one that would slide onto the finger all the way to the base without causing friction against the skin, using the same operationalization of friction as given in the instructions to participants above. Size was quantified as the inner diameter of the ring gauge.

## *Analysis*

For each participant and each finger, the thresholds from the two staircases were averaged. Percent overestimation finger size was calculated as  $100 * (\text{judged diameter} - \text{actual diameter}) / \text{actual diameter}$ , as in previous studies of distorted hand representation (e.g., Longo & Haggard, 2010, 2012a, 2012b; Longo et al., 2015; Tamè et al., 2017). Positive values of this measure indicate overestimation of finger size, whereas negative values indicate underestimation. We quantified ring gauge size in terms of the diameter of the ring, rather than the circumference. However, as the circumference is equal to the diameter multiplied by a constant value (i.e.,  $\pi$ ), these measures are mathematically equivalent. Note also that percent overestimation is mathematically equivalent to the body perception index (BPI), which has been widely used in body size estimation studies of patients with eating disorders (e.g., Slade & Russell, 1973; Pierloot & Houben, 1978; Thompson, Berland, Linton, & Weinsier, 1986; for discussion see Smeets, Smit, Panhuysen, & Ingleby, 1998), but defines veridical judgments with a value of 0, rather than 100 as in the BPI.


The preceding analysis quantifies finger size in terms of the diameter of the ring gauge, although the participant's task was not to judge 1-dimensional diameter. We therefore conducted a second analysis expressing the size of the ring gauges, and of the participant's finger, in terms of 2-dimensional area. Each threshold was re-expressed as an area (i.e.,  $\pi * (\text{diameter}/2)^2$ ), as was the size of each finger (note that this assumes that the cross-section of each finger is circular, which is only approximately true). Because the area increases as the square of the linear dimensions, it is logically possible for qualitatively different patterns to hold for these two analyses.

For both dependent measures, repeated-measures analysis of variance (ANOVA) was used to test for differences in overestimation of size across fingers. Where

Mauchley's test indicated a violation of the sphericity assumption, the Greenhouse-Geisser correction was applied. For each finger, one-sample t-tests were used to compare judgments to veridical performance.


### *Results and Discussion*

We first investigated the relation between actual and judged finger size. Figure 2 shows scatterplots of the relation between actual and judged diameter (top row) and area (bottom row). Robust correlations were apparent for both measures in all five fingers (all  $p$ 's < 0.002). These results demonstrate both that participants were effectively able to perform the task and also that their judgments assessed their own self-specific body image, rather than a generic representation of fingers in general. These correlations are also notable in light of the fact that other measures of body size estimation (e.g., the moving caliper method) have been criticized on the basis that judged body size did not correlate with actual size (e.g., Ben-Tovim & Crisp, 1984; Ben-Tovim, Walker, Murray, & Chin, 1990).



**Figure 2:** The relation between actual and judged finger size. *Top row:* scatterplots showing the relation between actual and judged ring diameter, for each of the five fingers. *Bottom row:* the same data for ring area. Grey lines show the least-squares regression line. Clear correlations between judged and actual finger size were apparent for all fingers.

The left panel of Figure 3 shows the results expressed in terms of the diameter of the rings. The magnitude of overestimation differed significantly across the five fingers,  $F(4, 76) = 3.35, p < 0.02, \eta_p^2 = 0.15$ . Absolute underestimation was found for the non-thumb fingers, though this only reached significance for the index finger ( $M: 5.51\%$  underestimation),  $t(19) = -3.49, p < 0.005, d = 0.78$ , and the ring finger ( $M: 4.16\%$  underestimation),  $t(19) = -2.22, p < 0.05, d = 0.50$ . No significant biases were found for the thumb ( $M: 1.00\%$  overestimation),  $t(19) = 0.53, p < 0.20, d = 0.12$ , the ring finger ( $M: 3.77\%$  underestimation),  $t(19) = -1.81, p = 0.086, d = 0.40$ , or the little finger ( $M: 1.43\%$  underestimation),  $t(19) = -0.51, p < 0.20, d = 0.11$ .



**Figure 3:** Results from Experiment 1. *Left panel:* overestimation of finger size as quantified by the diameter of the rings. *Right panel:* overestimation of finger size as quantified by the area of the rings. Error bars are one S.E.M. Positive values indicate overestimation of finger size, while negative values indicate underestimation.

The right panel of Figure 3 shows the same data expressed in terms of the circular area of the rings. Results were similar to those above, with a significant difference in the magnitude of overestimation across fingers,  $F(4, 76) = 3.43, p < 0.05, \eta_p^2 = 0.15$ . There was significant underestimation of the index finger ( $M: 10.25\%$  underestimation),  $t(19) = -3.51, p < 0.005, d = 0.78$ , and marginally significant

underestimation of the middle finger ( $M$ : 7.48 % underestimation),  $t(19) = -2.06$ ,  $p = 0.053$ ,  $d = 0.46$ . There were no significant biases for the thumb ( $M$ : 2.71 % overestimation),  $t(19) = 0.69$ ,  $p < 0.20$ ,  $d = 0.15$ , the ring finger ( $M$ : 6.57 % underestimation),  $t(19) = -1.58$ ,  $p = 0.131$ ,  $d = 0.35$ , or the little finger ( $M$ : 1.34 % underestimation),  $t(19) = -0.24$ ,  $p < 0.20$ ,  $d = 0.05$ .

## Experiment 2

The results of Experiment 1 provide evidence for systematic distortions of perceived finger size in 3-D. In Experiment 2, we tested whether these distortions were affected by the actual configuration adopted by the fingers. Several studies have demonstrated that the internal posture of the hands produces a range of changes in tactile processing both perceptually (e.g., Medina & Rapp, 2008; Overvliet et al., 2011; Riemer et al., 2010; Tamè et al., 2011, 2017b; Zampini et al., 2005) and neurally (e.g., Hamada & Suzuki, 2003, 2005; Sakata, Takaoka, Kawarasaki, & Shibutani, 1973; Stavrinou et al., 2007). Moreover, recent studies have found that changes in the internal posture of the hand (with fingers splayed vs pressed together) modulate the perceived distance between touches on the hand (Longo, 2017b) as well as proprioceptive perceptual hand maps (Longo, 2015b). It is unclear whether such changes in hand posture also affect the conscious body image. Here, we therefore tested whether posture modulates the internal representation of finger size.

## *Method*

### *Participants*

Twenty individuals (8 women, mean age = 29.7 years, range: 19-48 years) participated. Participants were generally right-handed as assessed by the Edinburgh

Inventory (mean = 85.4, range = -29.4 – 100). Participants reported no known abnormalities of tactile perception and normal or corrected-to-normal visual acuity.

### *Procedure*

Stimuli and procedures were identical to Experiment 1, except that stimuli were presented while the participant held their left hand in two different postures, as shown in Figure 4. In the *Together* posture, the participant held their left hand on the tabletop with the fingers pressed together. In the *Apart* posture, the participant held their hand with the fingers splayed as far apart as would be comfortable to maintain throughout each experimental block. As in Experiment 1, the participant was not allowed to see their hand at any time during the experiment, and it was covered with a black cloth.

The structure of each block was identical to Experiment 1, but there were now 10 blocks, one for each finger in each posture. The order of the 10 blocks was randomized. To check that the participant was following the instructions, the experimenter took a photograph of the participant's hand just before and after each block and before each break. Participants were required to respond based on the size of the fingers of their left hand.



**Figure 4:** The two postures in Experiment 2. *Left panel:* the *Together* posture, in which the hand was held with the fingers pressed together. *Right panel:* the *Apart* posture, in which the hand was held with the fingers splayed at the maximum amount that would be comfortable to hold throughout the experimental block.


### *Analysis*

Data analysis was identical to Experiment 1 except that posture was added as an additional factor to ANOVAs. In addition, we used Bayesian repeated-measures ANOVA (Wetzels, Grasman, & Wagenmakers, 2012), as implemented in JASP version 0.8.1.1 (JASP Team, 2018), to investigate whether the absence of a significant interaction between finger and posture provided positive evidence for the absence of an effect.

### *Results and Discussion*

Figure 5 shows the relation between actual and judged finger size, collapsed across the two postures. As in Experiment 1, positive correlations were apparent in all cases, though a few of them were somewhat weaker than in the previous experiment.






**Figure 5:** The relation between actual and judged finger size, collapsed across the two postures. *Top row:* scatterplots showing the relation between actual and judged ring diameter, for each of the five fingers. *Bottom row:* the same data for ring areas. Grey lines show the least-squares regression line. Positive correlations were apparent in all cases, though not all of them reached statistical significance.

The left panel of Figure 6 shows overestimation of finger size expressed in terms of ring diameter. As in Experiment 1, the magnitude of overestimation differed significantly across the fingers,  $F(2.55, 48.48) = 7.57, p < 0.0001, \eta_p^2 = 0.29$ . There was no significant effect of posture,  $F(1, 19) = 0.27, p > 0.20, \eta_p^2 = 0.01$ , nor an interaction of posture and finger,  $F(3.34, 63.37) = 1.33, p > 0.10, \eta_p^2 = 0.07$ . To investigate whether the data provide positive evidence against the presence of a postural effect, we conducted a Bayesian repeated-measures ANOVA. The Bayes Factor for the interaction provided substantial support for the null hypothesis,  $BF_{01} = 9.87$ .

Collapsing across the two postures, there was significant underestimation of the index finger ( $M: 9.51\%$  underestimation),  $t(19) = -4.98, p < 0.0001, d = 1.11$ , the middle finger ( $M: 6.61\%$  underestimation),  $t(19) = -2.60, p < 0.02, d = 0.58$ , and the ring finger ( $M: 5.61\%$  underestimation),  $t(19) = -2.50, p < 0.05, d = 0.55$ . As in Experiment 1, there were no significant biases for the thumb ( $M: 0.96\%$  underestimation),  $t(19) = -0.46, p > 0.20, d = 0.10$ , or the little finger ( $M: 3.54\%$  underestimation),  $t(19) = -1.28, p > 0.20, d = 0.29$ .



**Figure 6:** Results from Experiment 2. *Left panel:* overestimation of finger size as quantified by the diameter of the rings. *Right panel:* overestimation of finger size as quantified by the area of the rings. Error bars are one S.E.M. Positive values indicate overestimation of finger size, while negative values indicate underestimation.

The right panel of Figure 6 shows data from Experiment 2 expressed in terms of the circular area of rings. There was a significant difference in the magnitude of overestimation across fingers,  $F(2.52, 47.80) = 7.30, p < 0.001, \eta_p^2 = 0.28$ . However, there was no significant effect of posture,  $F(1, 19) = 0.45, p > 0.20, \eta_p^2 = 0.02$ , nor an interaction of posture and finger,  $F(3.30, 62.62) = 1.47, p > 0.20, \eta_p^2 = 0.07$ . A Bayesian repeated-measures ANOVA provided substantial support for the absence of an interaction of finger and posture,  $BF_{01} = 9.50$ .

Collapsing across the two postures, there was significant underestimation of the index finger ( $M: 17.29\%$  underestimation),  $t(19) = -4.97, p < 0.0001, d = 1.11$ , the middle finger ( $M: 11.48\%$  underestimation),  $t(19) = -2.36, p < 0.05, d = 0.53$ , and the ring finger ( $M: 9.84\%$  underestimation),  $t(19) = -2.26, p < 0.05, d = 0.51$ . There were no significant biases for the thumb ( $M: 0.98\%$  underestimation),  $t(19) = -0.23, p > 0.20, d = 0.05$ , or the little finger ( $M: 5.37\%$  underestimation),  $t(19) = -1.00, p > 0.20, d = 0.22$ .

## General Discussion

These results demonstrate systematic biases in the perceived 3-D size of the fingers. In both experiments, clear underestimation of the index finger (and to a lesser extent the middle finger) was present, which decreased progressively across the hand to the little finger. These biases were not affected by changes in the internal posture of the hand. Importantly, this pattern is qualitatively different from the pattern of underestimation found for 1-D judgments of finger length (Longo & Haggard, 2012a; Longo et al., 2015b; Tamè et al., 2017a). This suggests that judgments of the body as a coherent 3-D object may differ in fundamental ways from judgments about individual body dimensions.

As in the present study, those studies reported overall underestimation of the size of the non-thumb fingers. In contrast, however, the pattern across the four fingers was completely reversed. In the studies of perceived finger length, the magnitude of underestimation increased progressively from the index finger to the little finger, with the largest distortion found for the ring and little finger. This pattern mirrors the cortical magnification of each finger in somatosensory cortex (Duncan & Boynton, 2007) and the spatial sensitivity of the fingers (Duncan & Boynton, 2007; Manser-Smith et al., 2018; Vega-Bermudez & Johnson, 2001), both of which decrease progressively from the index finger to the little finger, as well as the increase in underestimation of finger length in proprioceptive hand maps (e.g., Longo & Haggard, 2010; Ganea & Longo, 2017; Cocchini, Galligan, Mora, & Kuhn, 2018). In contrast, in the present study, the largest underestimation was found for the index finger, and underestimation decreased across the hand towards the little finger. The exact causes of these changes are unclear. However, this pattern suggests that judgments of the fingers in 3-D are importantly different from 1-D judgments of length. This is consistent with the recent results of

Sadibolova and colleagues (2019) who found that judgments of body part volume showed a qualitatively different pattern of distortions from judgments of length.

One potential question about the present study is whether our task really required participants to make judgments of their fingers as 3-D objects. One might note that our actual quantifications are of ring diameter (a 1-D property) and area (a 2-D property), and not of an intrinsically 3-D measure such as volume. In our view, what makes this task assess the 3-D image of the fingers is the fact that participants have to imagine the finger going *through* the ring, and therefore need to consider the circumference of each finger along its entire length. The task used in the current study is interestingly different from that used by Sadibolova and colleagues (2019), who asked people to make explicit estimates of how many multiples of the volume of their hand or of a non-body object would make up the volume of different parts of their body. While we have argued that the current task does assess the participant's experience of their finger as a volumetric 3-D object, it clearly does not measure perceived finger volume. While this does make it somewhat more ambiguous exactly what aspect of body image we are measuring in the present study compared to the study of Sadibolova and colleagues (2019), it is worth emphasising that making quantitative judgments of how many multiples of the volume of an object make up the volume of a specific body part is a highly novel and non-intuitive judgment for participants to make. In contrast, judging whether a ring will fit on one's finger is a familiar judgment to make, which relates much more closely to people's everyday experiences and activities.

In this respect, the task used in this study has similarities to studies investigating judgments of affordances, such as the ability to pass through apertures (e.g., Warren & Whang, 1987; Graydon, Linkenauger, Teachman, & Proffitt, 2012), to reach objects (e.g., Carello et al., 1989; Rochat & Wraga, 1997), and to sit on stools (e.g., Mark, 1987).

Because we asked participants about the ability of part of their body to fit through a form of aperture, there is a sense in which our task is also an affordance judgment (cf. Gibson, 1979). The potential link to affordances, however, is particularly intriguing in light of several recent studies which have found that judgments of the ability to pass through apertures are altered in individuals with anorexia nervosa (e.g., Guardia et al., 2010, 2012; Keizer et al., 2013; Metral et al., 2014) and in healthy adults following embodiment of avatars of different body sizes in virtual reality (Piryankova et al., 2014). Such results suggest that the implicit use of body size information in making affordance judgments has functional connections with the conscious body image. A recent study by Engel and Keizer (2017) is especially intriguing in that it used a task analogous to ours in which participants had to estimate whether hoops of different diameter would fit around their entire body. Patients with anorexia overestimated the required hoop size compared to control participants. Subsequently, Keizer and colleagues (in press) found that 8-weeks of training with this task in patients with anorexia led to improvements in both affordance judgments and tactile distortions.

One recent study found that changes in perceived finger size following cutaneous anaesthesia (cf. Gandevia & Phegan, 1999) specifically affect judgments of finger width but not finger length (Walsh et al., 2015). Another study by Hashimoto and Iriki (2013) identified distinct patterns of brain activation using fMRI when participants made judgments about different body axes. Such effects are intriguing as they suggest that there may be distinct mental representations of individual body axes. As mentioned in the previous paragraph, however, this does not imply that the experience of the body as a 3-D object can be predicted from judgments of each body axis in isolation. Rather 3-D perception of the body may rely on distinct mental representations from judgments about individual body dimensions. It is intriguing in this light that whereas 1-D

judgments of finger length appear to mirror differences between the fingers in sensitivity and cortical magnification factors (Longo & Haggard, 2012a), this relation does not hold for judgments of 3-D finger size in the present study.

This study contributes to an emerging literature detailing the way in which the body is represented as a volumetric, 3-D object in the world. One set of studies has approached this issue by comparing perception of the palmar and dorsal surfaces of the hand. These skin surfaces have different physiological properties (Mountcastle, 2005) and distinct representations in somatosensory cortex (Merzenich, Kaas, Sur, & Lin, 1978; Nelson, Sur, Felleman, & Kaas, 1980). Perceptual effects arising from low-level sensory maps of the hand may therefore show highly distinct patterns of performance on each surface. In contrast, if perceptual effects arise from higher-level representations of the body as a volumetric whole, similar patterns should be found on each surface. Interestingly, perceptual abilities appear to differ widely in this regard. For example, tactile localisation (e.g., Mancini et al., 2011) and tactile distance perception (e.g., Longo & Haggard, 2011; Longo, Ghosh, & Yahya, 2015) appear to arise from fragmented 2-D maps of individual skin surfaces, as the pattern of distortions on each surface (i.e., palm and dorsum) differs. Tactile confusions between fingers (Manser-Smith et al., 2018) and explicit judgments of hand shape (Longo, 2015d), on the other hand, appear to arise from coherent 3-D representations of whole body parts, as the two sides of the hand show highly similar patterns of perceptual bias.

There was no apparent effect of posture on judgments of finger size. This is in marked contrast to a number of recent studies that have found that finger and arm posture modulates tactile processing (e.g., Longo, 2017b; Medina & Rapp, 2008; Overvliet et al., 2011; Riemer et al., 2010; Sadibolova, Tamè, & Longo, 2018; Tamè et al., 2011, 2017b; Zampini et al., 2005), proprioceptive body maps (Longo, 2015b), and

somatotopic maps in somatosensory cortex (Hamada & Suzuki, 2003, 2005; Stavrinou et al., 2007). This pattern suggests that splaying the fingers may modulate low-level somatosensory representations of the body, composed of fragmented 2-D maps of individual skin surfaces, but not higher-level representations of the body as a 3-D volumetric object.

It is noteworthy that judgments of finger size in the present study were strongly correlated with actual finger size. Some authors have highlighted the lack of such correlations for tasks such as the moving caliper method, arguing that this calls into question the extent to which distortions using this method are actually perceptual, rather than attitudinal (e.g., Ben-Tovim et al., 1979, 1990; Ben-Tovim & Crisp, 1984; see also Smeets, Smit, Panhuysen, & Ingleby, 1998). The presence of clear correlations in this study is therefore an important validation that participants are in fact basing their judgments on their experience of their own fingers specifically, rather than a more generic representations of fingers in general. It further provides support for the interpretation that the distortions we report are genuinely perceptual distortions, and not artefacts of attitudes towards the fingers.

The present results contribute to a growing literature showing a range of systematic misperception and distortions of the mental representation of the hand (for review, see Longo, 2017a), focusing on the body as a three-dimensional object. Large and highly stereotyped distortions have been found for perceptual abilities including tactile localisation (e.g., Mancini et al., 2011; Medina, Tamè, & Longo, 2018), tactile distance perception (e.g., Taylor-Clarke et al., 2004; Fiori & Longo, 2018), and position sense (e.g., Longo & Haggard, 2010, 2012b), as well as more explicit judgments of the location of landmarks within the hand (e.g., Longo, 2015a; Ambroziak et al., 2018), hand proportions (e.g., Longo & Haggard, 2012a; Longo & Sadibolova, 2013) and finger size

(this study, Longo & Haggard, 2012a; Tamè et al., 2017a). Despite the hand's role as a paragon of familiarity and intimate knowledge, the brain nevertheless maintains highly distorted representations of the hand, which have widespread influences on perception.



## References

- Allebeck, P., Hallberg, D., & Espmark, S. (1976). Body image – An apparatus for measuring disturbances in estimation of size and shape. *Journal of Psychosomatic Research*, 20, 583-589.
- Ambroziak, K. B., Tamè, L., & Longo, M. R. (2018). Conceptual distortions of hand structure are robust to changes in stimulus information. *Consciousness and Cognition*, 61, 107-116.
- Ben-Tovim, D. I., & Crisp, A. H. (1984). The reliability of estimates of body width and their relationship to current measured body size among anorexic and normal subjects. *Psychological Medicine*, 14, 843-846.
- Ben-Tovim, D. I., Walker, M. K., Murray, H., & Chin, G. (1990). Body size estimates: Body image or body attitude measure? *International Journal of Eating Disorders*, 9, 57-67.
- Ben-Tovim, D. I., Whitehead, J., & Crisp, A. H. (1979). A controlled study of the perception of body width in anorexia nervosa. *Journal of Psychosomatic Research*, 23, 267-272.
- Brainard, D. H. (1997). The psychophysics toolbox. *Spatial Vision*, 10, 433-436.
- Brodie, D. A., Slade, P. D., & Rose, H. (1989). Reliability measures in distorting body-image. *Perceptual and Motor Skills*, 69, 723-732.
- Bruch, H. (1962). Perceptual and conceptual disturbances in anorexia nervosa. *Psychosomatic Medicine*, 24, 187-194.
- Button, E. J., Fransella, F., & Slade, P. D. (1977). A reappraisal of body perception disturbance in anorexia nervosa. *Psychological Medicine*, 7, 235-243.
- Carello, C., Groszofsky, A., Reichel, F. D., Solomon, H. Y., & Turvey, M. T. (1989). Visually perceiving what is reachable. *Ecological Psychology*, 1, 25-54.

- Cash, T. F., & Deagle, E. A. (1997). The nature and extent of body-image disturbances in anorexia nervosa and bulimia nervosa: A meta-analysis. *International Journal of Eating Disorders*, 22, 107-125.
- Casper, R. C., Halmi, K. A., Goldberg, S. C., Eckert, E. D., & Davis, J. M. (1979). Disturbances in body image estimation as related to other characteristics and outcome in anorexia nervosa. *British Journal of Psychiatry*, 134, 60-66.
- Cocchini, G., Galligan, T., Mora, L., & Kuhn, G. (2018). The magic hand: Plasticity of mental hand representation. *Quarterly Journal of Experimental Psychology*, 71, 2314-2324.
- Coelho, L. A., Zaninelli, G., & Gonzalez, C. L. R. (2017). A kinematic examination of hand perception. *Psychological Research*, 81, 1224-1231.
- D'Amour, S., & Harris, L. R. (2017). Perceived face size in healthy adults. *PLOS ONE*, 12, e0177349.
- Dolan, B. M., Birtchnell, S. A., & Lacey, J. H. (1987). Body image distortion in non-eating disordered women and men. *Journal of Psychosomatic Research*, 31, 513-520.
- Dolce, J. J., Thompson, J. K., Register, A., & Spana, R. E. (1987). Generalization of body size distortion. *International Journal of Eating Disorders*, 6, 401-408.
- Duncan, R. O., & Boynton, G. M. (2007). Tactile hyperacuity thresholds correlate with finger maps in primary somatosensory cortex (S1). *Cerebral Cortex*, 17, 2878-2891.
- Engel, M. M., & Keizer, A. (2017). Body representation disturbances in visual perception and affordance perception persist in eating disorder patients after completing treatment. *Scientific Reports*, 7, 16184.
- Fernández, F., Probst, M., Meermann, R., & Vandereycken, W. (1994). Body size

- estimation and body dissatisfaction in eating disorder patients and normal controls. *International Journal of Eating Disorders*, 16, 307-310.
- Fiori, F., & Longo, M. R. (2018). Tactile distance illusions reflect a coherent stretch of tactile space. *Proceedings of the National Academy of Sciences*, 115, 1238-1243.
- Freeman, R. J., Thomas, C. D., Solyom, L., & Hunter, M. A. (1984). A modified video camera for measuring body image distortion: Technical description and reliability. *Psychological Medicine*, 14, 411-416.
- Fuentes, C. T., Longo, M. R., & Haggard, P. (2013). Body image distortions in healthy adults. *Acta Psychologica*, 144, 344-351.
- Gandevia, S. C., & Phegan, C. M. L. (1999). Perceptual distortions of the human body image produced by local anaesthesia, pain and cutaneous stimulation. *Journal of Physiology*, 514.2, 609-616.
- Ganea, N., & Longo, M. R. (2017). Projecting the self outside the body: Body representations underlying proprioceptive imagery. *Cognition*, 162, 41-47.
- Garner, D. M., Garfinkel, P. E., Stancer, H. C., & Moldofsky, H. (1976). Body image disturbances in anorexia nervosa and obesity. *Psychosomatic Medicine*, 38, 327-336.
- Gibson, J. J. (1979). *The ecological approach to visual perception*. Boston, MA: Houghton Mifflin.
- Glucksman, M. L., & Hirsch, J. (1969). The response of obese patients to weight reduction: III. The perception of body size. *Psychosomatic Medicine*, 31, 1-7.
- Graydon, M. M., Linkenauger, S. A., Teachman, B. A., & Proffitt, D. R. (2012). Scared stiff: The influence of anxiety on the perception of action capabilities. *Cognition & Emotion*, 26, 1301-1315.
- Green, B. G. (1982). The perception of distance and location for dual tactile pressures.

- Perception and Psychophysics*, 31, 315-323.
- Guardia, D., Lafargue, G., Thomas, P., Dodin, V., Cottencin, O., & Luyat, M. (2010). Anticipation of body-scaled action is modified in anorexia nervosa. *Neuropsychologia*, 48, 3961-3966.
- Guardia, D., Conversy, L., Jardri, R., Lafargue, G., Thomas, P., Dodin, V., Cottencin, O., & Luyat, M. (2012). Imagining one's own and someone else's body actions: Dissociation in anorexia nervosa. *PLOS ONE*, 7, e43241.
- Halmi, K. A., Goldberg, S. C., & Cunningham, S. (1977). Perceptual distortion of body image in adolescent girls: Distortion of body image in adolescence. *Psychological Medicine*, 7, 253-257.
- Hamada, Y., & Suzuki, R. (2003). Hand posture modulates neuronal interaction in the primary somatosensory cortex of humans. *Clinical Neurophysiology*, 114, 1689-1696.
- Hamada, Y., & Suzuki, R. (2005). Hand posture modulates cortical finger representation in SII. *NeuroImage*, 25, 708-717.
- Hashimoto, T., & Iriki, A. (2013). Dissociations between the horizontal and dorsoventral axes in body-size perception. *European Journal of Neuroscience*, 37, 1747-1753.
- Horne, R. L., Van Vactor, J. C., & Emerson, S. (1991). Disturbed body image in patients with eating disorders. *American Journal of Psychiatry*, 148, 211-215.
- Hundleby, J. D., & Bourguoin, N. C. (1993). Generality in the errors of estimation of body image. *International Journal of Eating Disorders*, 13, 85-92.
- JASP Team. (2018). JASP (Version 0.8.1.1).
- Jeannerod, M. (1997). *The cognitive neuroscience of action*. Oxford: Blackwell.
- Keizer, A., Engel, M. M., Bonekamp, J., & van Elburg, A. (in press). Hoop training: A pilot

- study assessing the effectiveness of a multisensory approach to treatment of body image disturbance in anorexia nervosa. *Eating and Weight Disorders*.
- Keizer, A., Smeets, M. A. M., Dijkerman, H. C., Uzunbajakou, S. A., van Elburg, A., & Postma, A. (2013). Too fat to fit through the door: First evidence for disturbed body-scaled action in anorexia nervosa. *PLOS ONE*, *8*, e64602.
- Keizer, A., van Elburg, A., Helms, R., & Dijkerman, H. C. (2016). A virtual reality full body illusion improves body image disturbance in anorexia nervosa. *PLOS ONE*, *11*, e0163921.
- Lindholm, L., & Wilson, G. T. (1988). Body image assessment in patients with bulimia nervosa and normal controls. *International Journal of Eating Disorders*, *7*, 527-539.
- Linkenauger, S. A., Wong, H. Y., Geuss, M., Stefanucci, J. K., McCulloch, K. C., Bühlhoff, H. H., et al. (2015). The perceptual homunculus: The perception of the relative proportions of the human body. *Journal of Experimental Psychology: General*, *144*, 103-113.
- Linkenauger, S. A., Kirby, L. R., McCulloch, K. C., & Longo, M. R. (2017). People watching: The perception of the relative body proportions of the self and others. *Cortex*, *92*, 1-7.
- Longo, M. R. (2015a). Intuitive anatomy: Distortions of conceptual knowledge of hand structure. *Cognition*, *142*, 230-235.
- Longo, M. R. (2015b). Posture modulates implicit hand maps. *Consciousness and Cognition*, *36*, 96-102.
- Longo, M. R. (2015c). Implicit and explicit body representations. *European Psychologist*, *20*, 6-15.
- Longo, M. R. (2015d). Three-dimensional coherence of the conscious body

- image. *Quarterly Journal of Experimental Psychology*, 68, 1116-1123.
- Longo, M. R. (2017a). Distorted body representations in healthy cognition. *Quarterly Journal of Experimental Psychology*, 70, 378-388.
- Longo, M. R. (2017b). Hand posture modulates perceived tactile distance. *Scientific Reports*, 7, 9665.
- Longo, M. R., & Haggard, P. (2010). An implicit body representation underlying human position sense. *Proceedings of the National Academy of Sciences*, 107, 11727-11732.
- Longo, M. R., & Haggard, P. (2011). Weber's illusion and body shape: Anisotropy of tactile size perception on the hand. *Journal of Experimental Psychology: Human Perception and Performance*, 37, 720-726.
- Longo, M. R., & Haggard, P. (2012a). Implicit body representations and the conscious body image. *Acta Psychologica*, 141, 164-168.
- Longo, M. R., & Haggard, P. (2012b). A 2.5-D representation of the human hand. *Journal of Experimental Psychology: Human Perception and Performance*, 38, 9-13.
- Longo, M. R., & Sadibolova, R. (2013). Seeing the body distorts tactile size perception. *Cognition*, 126, 475-481.
- Longo, M. R., Ghosh, A., & Yahya, T. (2015). Bilateral symmetry of distortions of tactile size perception. *Perception*, 44, 1251-1262.
- Longo, M. R., Long, C., & Haggard, P. (2012). Mapping the invisible hand: A body model of a phantom limb. *Psychological Science*, 23, 740-742.
- Longo, M. R., Mancini, F., & Haggard, P. (2015a). Implicit body representations and tactile spatial remapping. *Acta Psychologica*, 160, 77-87.
- Longo, M. R., Mattioni, S., & Ganea, N. (2015b). Perceptual and conceptual distortions of implicit hand maps. *Frontiers in Human Neuroscience*, 9, 656.

- Mancini, F., Longo, M. R., Iannetti, G. D., & Haggard, P. (2011). A supramodal representation of the body surface. *Neuropsychologia*, 49, 1194-1201.
- Manser-Smith, K., Tamè, L., & Longo, M. R. (2018). Tactile confusions of the fingers and toes. *Journal of Experimental Psychology: Human Perception and Performance*, 44, 1727-1738.
- Margolis, A. N., & Longo, M. R. (2015). Visual detail about the body modulates tactile localisation biases. *Experimental Brain Research*, 233, 351-358.
- Mark, L. S. (1987). Eyeheight-scaled information about affordances: A study of sitting and stair climbing. *Journal of Experimental Psychology: Human Perception and Performance*, 13, 361-370.
- Medina, J., & Rapp, B. (2008). Phantom tactile sensations modulated by body position. *Current Biology*, 18, 1937-1942.
- Medina, S., Tamè, L., & Longo, M. R. (2018). Tactile localization biases are modulated by gaze direction. *Experimental Brain Research*, 236, 31-42.
- Merzenich, M. M., Kaas, J. H., Sur, M., & Lin, C.-S. (1978). Double representation of the body surface within cytoarchitecture areas 3b and 1 in "S1" in the owl monkey (*aotus trivirgatus*). *Journal of Comparative Neurology*, 181, 41-73.
- Metral, M., Guardia, D., Bauwens, I., Guerraz, M., Lafargue, G., Cottencin, O., & Luyat, M. (2014). Painfully thin but locked inside a fatter body: Abnormalities in both anticipation and execution of action in anorexia nervosa. *BMC Research Notes*, 7, 707.
- Mölbart, S. C., Klein, L., Thaler, A., Mohler, B. J., Brozzo, C., ..., & Giel, K. E. (2017). Depictive and metric body size estimation in anorexia nervosa and bulimia nervosa: A systematic review and meta-analysis. *Clinical Psychology Review*, 57, 21-31.

- Mölbart, S. C., Sauer, H., Dammann, D., Zipfel, S., Teufel, M., ... & Mack, I. (2016). Multimodal body representation of obese children and adolescents before and after weight-loss treatment in comparison to normal-weight children. *PLOS ONE*, *11*, e0166826.
- Mountcastle, V. B. (2005). *The sensory hand: Neural mechanisms of somatic sensation*. Cambridge, MA: Harvard University Press.
- Nelson, R. J., Sur, M., Felleman, D. J., & Kaas, J. H. (1980). Representations of the body surface in postcentral parietal cortex of macaca fascicularis. *Journal of Comparative Neurology*, *192*, 611–643.
- Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia*, *9*, 97-113.
- Overvliet, K. E., Anema, H. A., Brenner, E., Dijkerman, H. C., & Smeets, J. B. (2011). Relative finger position influences whether you can localize tactile stimuli. *Experimental Brain Research*, *208*, 245-255.
- Pierloot, R. A., & Houben, M. E. (1978). Estimation of body dimensions in anorexia nervosa. *Psychological Medicine*, *8*, 317-324.
- Piryankova, I., Wong, H. Y., Linkenauger, S. A., Stinson, C., Longo, M. R., Bülthoff, H. H., & Mohler, B. J. (2014). Owning an overweight or underweight body: Distinguishing the physical, experienced and virtual body. *PLOS ONE*, *9*, e103428.
- Riemer, M., Trojan, J., Kleinböhl, D., & Hölzl, R. (2010). Body posture affects tactile discrimination and identification of fingers and hands. *Experimental Brain Research*, *206*, 47-57.
- Rochat, P., & Wraga, M. (1997). An account of the systematic error in judging what is reachable. *Journal of Experimental Psychology: Human Perception and Performance*, *23*, 199-212.



- Ruff, G. A., & Barrios, B. A. (1986). Realistic assessment of body image. *Behavioral Assessment, 8*, 237-251.
- Sadibolova, R., Ferrè, E. R., Linkenauger, S. A., & Longo, M. R. (2019). Distortions of perceived volume and length of body parts. *Cortex, 111*, 74-86.
- Sadibolova, R., Tamè, L., & Longo, M. R. (2018). More than skin-deep: Integration of skin-based and musculo-skeletal reference frames in localisation of touch. *Journal of Experimental Psychology: Human Perception and Performance, 44*, 1672-1682.
- Sakata, H., Takaoka, Y., Kawarasaki, A., & Shibutani, H. (1973). Somatosensory properties of neurons in the superior parietal cortex (area 5) of the rhesus monkey. *Brain Research, 64*, 85-102.
- Salbach, H., Klinkowski, N., Pfeiffer, E., Lehmkuhl, U., & Korte, A. (2007). Body image and attitudinal aspects of eating disorders in rhythmic gymnasts. *Psychopathology, 40*, 388-393.
- Sanabria, D., Soto-Faraco, S., & Spence, C. (2005). Spatiotemporal interactions between audition and touch depend on hand posture. *Experimental Brain Research, 165*, 505-514.
- Sathian, K., & Zangaladze, A. (1996). Tactile spatial acuity at the human fingertip and lip: Bilateral symmetry and inter-digit variability. *Neurology, 46*, 1464-1466.
- Saulton, A., Dodds, T. J., Bulthöff, H. H., & de la Rosa, S. (2015). Objects exhibit body model like distortions. *Experimental Brain Research, 233*, 1471-1499.
- Schneider, N., Frieler, K., Pfeiffer, E., Lehmkuhl, U., & Salbach-Andrae, H. (2009a). Comparison of body size estimation in adolescents with different types of eating disorders. *European Eating Disorders Review, 17*, 468-475.
- Schneider, N., Martus, P., Ehrlich, S., Pfeiffer, E., Lehmkuhl, U., & Salbach-Andrae, H.

- (2009b). The assessment of body image distortion in female adolescents with anorexia nervosa: The development of a test for body image distortion in children and adolescents (BID-CA). *Eating and Weight Disorders*, 14, e128-e136.
- Sepúlveda, A. R., Botella, J., & León, J. A. (2002). Body-image disturbance in eating disorders: A meta-analysis. *Psychology in Spain*, 6, 83-95.
- Slade, P. D., & Russell, G. F. M. (1973). Awareness of body dimensions in anorexia nervosa: Cross-sectional and longitudinal studies. *Psychological Medicine*, 3, 188-199.
- Smeets, M. A. M., Smit, F., Panhuysen, G. E. M., & Ingleby, J. D. (1997). The influence of methodological differences on the outcome of body size estimation studies in anorexia nervosa. *British Journal of Clinical Psychology*, 36, 263-277.
- Smeets, M. A. M., Smit, F., Panhuysen, G. E. M., & Ingleby, J. D. (1998). Body perception index: Benefits, pitfalls, ideas. *Journal of Psychosomatic Research*, 44, 457-464.
- Stavrinou, M. L., Della Penna, S., Pizzella, V., Torquati, K., Cianflone, F., et al (2007). Temporal dynamics of plastic changes in human primary somatosensory cortex after finger webbing. *Cerebral Cortex*, 17, 2134-2142.
- Tamè, L., Farnè, A., & Pavani, F. (2011). Spatial coding of touch at the fingers: Insights from double simultaneous stimulation within and between hands. *Neuroscience Letters*, 487, 78-82.
- Tamè, L., Bumpus, N., Linkenauger, S. A., & Longo, M. R. (2017). Distorted body representations are robust to differences in experimental instructions. *Attention, Perception, and Psychophysics*, 79, 1204-1216.
- Tamè, L., Dransfield, E., Quettier, T., & Longo, M. R. (2017). Finger posture modulates structural body representations. *Scientific Reports*, 7, 43019.
- Taylor-Clarke, M., Jacobsen, M., & Haggard, P. (2004). Keeping the world a constant size:

- Object constancy in human touch. *Nature Neuroscience*, 7, 219-220.
- Thompson, J. K. (1986). Larger than life. *Psychology Today*, 20, 38-44.
- Thompson, J. K., & Spana, R. E. (1988). The adjustable light beam method for the assessment of size estimation accuracy: Description, psychometric, and normative data. *International Journal of Eating Disorders*, 7, 521-526.
- Thompson, J. K., Berland, N. W., Linton, P. H., & Weinsier, R. L. (1986). Utilization of a self-adjusting light beam in the objective assessment of body distortion in seven eating disorder groups. *International Journal of Eating Disorders*, 5, 113-120.
- Traub, A. C., & Orbach, J. (1964). Psychophysical studies of body-image: I. The adjustable body-distorting mirror. *Archives of General Psychiatry*, 11, 53-66.
- Trojan, J., Kleinböhl, D., Stolle, A. M., Andersen, O. K., Hölzl, R., & Arendt-Nielsen, L. (2006). Psychophysical 'perceptual maps' of heat and pain sensations by direct localization of CO<sub>2</sub> laser stimuli on the skin. *Brain Research*, 1120, 106-113.
- Urdapilleta, I., Aspavlo, D., Masse, L., & Docteur, A. (2010). Use of a picture distortion technique to examine perceptive and ideal body image in male and female competitive swimmers. *Psychology of Sport and Exercise*, 11, 568-573.
- Vega-Bermudez, F., & Johnson, K. O. (2001). Differences in spatial acuity between digits. *Neurology*, 56, 1389-1391.
- Walsh, L. D., Hoad, D., Rothwell, J. C., Gandevia, S. C., & Haggard, P. (2015). Anesthesia changes perceived finger width but not finger length. *Experimental Brain Research*, 233, 1761-1771.
- Warren, W. H., & Whang, S. (1987). Visual guidance of walking through apertures: Body-scaled information for affordances. *Journal of Experimental Psychology: Human Perception and Performance*, 13, 371-383.
- Watson, A. B., & Pelli, D. G. (1983). QUEST: A Bayesian adaptive psychophysical method.

*Perception and Psychophysics*, 33, 113-120.

Wetzels, R., Grasman, R. P. P. P., & Wagenmakers, E.-J. (2012). A default Bayesian hypothesis test for ANOVA designs. *American Statistician*, 66, 104–111.

Yamamoto, S., & Kitazawa, S. (2001). Reversal of subjective temporal order due to arm crossing. *Nature Neuroscience*, 4, 759-765.

Zampini, M., Harris, C., & Spence, C. (2005). Effect of posture change on tactile perception: Impaired direction discrimination performance with interleaved fingers. *Experimental Brain Research*, 166, 498-508.

## Acknowledgments

This research was supported by an Erasmus Internship from the European Commission to EET and a grant from the European Research Council (ERC-2013-StG-336050) under the FP7 to MRL.